

Quality Measure Choices that Align with the Million Hearts® 2020 Initiative and Improving Cardiac Health
<https://qpp.cms.gov/mips/explore-measures> - Revised 09.23.19

Measure Name	Measure Description	eMEASURE ID	e MEASURE NQF	NQF	Quality ID	NQS Domain	Measure Trype	High Priority Measure	Data Submission Method	Speciality Measure Set	Primary Measure Steward
Controlling High Blood Pressure	Percentage of patients 18-85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled (<140/90mmHg) during the measurement period	CMS165v6	None	18	236	Effective Clinical Care	Intermediate Outcome	TRUE	Claims, CMS Web Interface, eCQMs	Cardiology, Obstetrics/Gynecology, Vascular Surgery, Thoracic Surgery, Rheumatology	National Committee for Quality Assurance
Coronary Artery Disease (CAD): Antiplatelet Therapy	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease (CAD) seen within a 12 month period who were prescribed aspirin or clopidogrel	None	None	67	6	Effective Clinical Care	Process	FALSE	CQMs	Cardiology,Family Medicine,Internal Medicine	American Heart Association
Ischemic Vascular Disease (IVD) All or None Outcome Measure (Optimal Control)	The IVD All-or-None Measure is one outcome measure (optimal control). The measure contains four goals. All four goals within a measure must be reached in order to meet that measure. The numerator for the all-or-none measure should be collected from the organization's total IVD denominator. All-or-None Outcome Measure (Optimal Control) - Using the IVD denominator optimal results include: ½ Most recent blood pressure (BP) measurement is less than 140/90 mm Hg -- And ½ Most recent tobacco status is Tobacco Free -- And ½ Daily Aspirin or Other Antiplatelet Unless Contraindicated -- And ½ Statin Use Unless Contraindicated	None	None	None	441	Effective Clinical Care	Intermediate Outcome	TRUE	CQMs	Cardiology,Family Medicine,Internal Medicine,Vascular Surgery,Thoracic Surgery	Wisconsin Collaborative for Healthcare Quality

Quality Measure Choices that Align with the Million Hearts® 2020 Initiative and Improving Cardiac Health
<https://qpp.cms.gov/mips/explore-measures> - Revised 09.23.19

Measure Name	Measure Description	eMEASURE ID	e MEASURE NQF	NQF	Quality ID	NQS Domain	Measure Type	High Priority Measure	Data Submission Method	Specialty Measure Set	Primary Measure Steward
Cardiac Rehabilitation Patient Referral from an Outpatient Setting	Percentage of patients evaluated in an outpatient setting who within the previous 12 months have experienced an acute myocardial infarction (MI), coronary artery bypass graft (CABG) surgery, a percutaneous coronary intervention (PCI), cardiac valve surgery, or cardiac transplantation, or who have chronic stable angina (CSA) and have not already participated in an early outpatient cardiac rehabilitation/secondary prevention (CR) program for the qualifying event/diagnosis who were referred to a CR program	None	None	643	243	Communication and Care Coordination	Process	TRUE	CQMs	Cardiology,Family Medicine,Internal Medicine	American College of Cardiology Foundation
Functional Status Assessments for Congestive Heart Failure	Percentage of patients 18 years of age and older with congestive heart failure who completed initial and follow-up patient-reported functional status assessments	CMS90v7	None	None	377	Person and Caregiver-Centered Experience and Outcomes	Process	TRUE	eCQMs	Family Medicine,Internal Medicine	Centers for Medicare & Medicaid Services
Adult Major Depressive Disorder (MDD): Coordination of Care of Patients with Specific Comorbid Conditions	Percentage of medical records of patients aged 18 years and older with a diagnosis of major depressive disorder (MDD) and a specific diagnosed comorbid condition (diabetes, coronary artery disease, ischemic stroke, intracranial hemorrhage, chronic kidney disease [stages 4 or 5], End Stage Renal Disease [ESRD] or congestive heart failure) being treated by another clinician with communication to the clinician treating the comorbid condition	None	None	None	325	Communication and Care Coordination	Process	TRUE	CQMs	Mental/Behavioral Health	American Psychiatric Association

Quality Measure Choices that Align with the Million Hearts® 2020 Initiative and Improving Cardiac Health
<https://qpp.cms.gov/mips/explore-measures> - Revised 09.23.19

Measure Name	Measure Description	eMEASURE ID	e MEASURE NQF	NQF	Quality ID	NQS Domain	Measure Type	High Priority Measure	Data Submission Method	Speciality Measure Set	Primary Measure Steward
Cardiac Stress Imaging Not Meeting Appropriate Use Criteria: Testing in Asymptomatic, Low-Risk Patients	Percentage of all stress single-photon emission computed tomography (SPECT) myocardial perfusion imaging (MPI), stress echocardiogram (ECHO), cardiac computed tomography angiography (CCTA), and cardiovascular magnetic resonance (CMR) performed in asymptomatic, low coronary heart disease (CHD) risk patients 18 years and older for initial detection and risk assessment	None	None	None	324	Efficiency and Cost Reduction	Efficiency	TRUE	CQMs	Cardiology	American College of Cardiology
Coronary Artery Disease (CAD): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy - Diabetes or Left Ventricular Systolic Dysfunction (LVEF < 40%)	Percentage of patients aged 18 years and older with a diagnosis of coronary artery disease seen within a 12 month period who also have diabetes OR a current or prior Left Ventricular Ejection Fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy	None	None	66	118	Effective Clinical Care	Process	FALSE	CQMs	Cardiology	American Heart Association
Heart Failure (HF): Angiotensin-Converting Enzyme (ACE) Inhibitor or Angiotensin Receptor Blocker (ARB) Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed ACE inhibitor or ARB therapy either within a 12 month period when seen in the outpatient setting OR at each hospital discharge	CMS135v6	2907	81	5	Effective Clinical Care	Process	FALSE	eCQMs	Cardiology,Family Medicine,Internal Medicine,Hospitalists	Physician Consortium for Performance Improvement

Quality Measure Choices that Align with the Million Hearts® 2020 Initiative and Improving Cardiac Health
<https://qpp.cms.gov/mips/explore-measures> - Revised 09.23.19

Measure Name	Measure Description	eMEASURE ID	e MEASURE NQF	NQF	Quality ID	NQS Domain	Measure Trype	High Priority Measure	Data Submission Method	Speciality Measure Set	Primary Measure Steward
Heart Failure (HF): Beta-Blocker Therapy for Left Ventricular Systolic Dysfunction (LVSD)	Percentage of patients aged 18 years and older with a diagnosis of heart failure (HF) with a current or prior left ventricular ejection fraction (LVEF) < 40% who were prescribed beta-blocker therapy either within a 12-month period when seen in the outpatient setting OR at each hospital discharge	CMS144v6	2908	83	8	Effective Clinical Care	Process	FALSE	eCQMs	Cardiology,Family Medicine,Internal Medicine,Hospitalists	Physician Consortium for Performance Improvement